



ICD-9-CM Code	ICD-10-CM Code(s)
<b>474.00</b>	Chronic tonsillitis
ICD-10-CM Codes	<p><b>J35.01</b> Chronic tonsillitis</p> <p>* Use additional code to identify:</p> <p>Exposure to environmental tobacco smoke (<b>Z77.22</b>)</p> <p>Exposure to tobacco smoke in the perinatal period (<b>P96.81</b>)</p> <p>History of tobacco use (<b>Z87.891</b>)</p> <p>Occupational exposure to environmental tobacco smoke (<b>Z57.31</b>)</p> <p>Tobacco dependence (<b>F17-</b>)</p> <p>Tobacco use (<b>Z72.0</b>)</p>
<b>382.3</b>	Unspecified chronic suppurative otitis media
ICD-10-CM Codes	<p><b>H66.3x1</b> Other chronic suppurative otitis media, right ear</p> <p><b>H66.3x2</b> Other chronic suppurative otitis media, left ear</p> <p><b>H66.3x3</b> Other chronic suppurative otitis media, bilateral</p> <p><b>H66.3x9</b> Other chronic suppurative otitis media, unspecified ear</p> <p>* Use additional code for any associated perforated tympanic membrane (<b>H72-</b>)</p> <p>* Use additional code to identify:</p> <p>Exposure to environmental tobacco smoke (<b>Z77.22</b>)</p> <p>Exposure to tobacco smoke in the perinatal period (<b>P96.81</b>)</p> <p>History of tobacco use (<b>Z87.891</b>)</p> <p>Occupational exposure to environmental tobacco smoke (<b>Z57.31</b>)</p> <p>Tobacco dependence (<b>F17-</b>)</p> <p>Tobacco use (<b>Z72.0</b>)</p>
<b>474.10</b>	Hypertrophy of tonsils with adenoids
ICD-10-CM Codes	<p><b>J35.3</b> Hypertrophy of tonsils with hypertrophy of adenoids</p> <p>* Use additional code to identify:</p> <p>Exposure to environmental tobacco smoke (<b>Z77.22</b>)</p> <p>Exposure to tobacco smoke in the perinatal period (<b>P96.81</b>)</p> <p>History of tobacco use (<b>Z87.891</b>)</p> <p>Occupational exposure to environmental tobacco smoke (<b>Z57.31</b>)</p> <p>Tobacco dependence (<b>F17-</b>)</p> <p>Tobacco use (<b>Z72.0</b>)</p>
<b>381.10</b>	Chronic serous otitis media, simple or unspecified
ICD-10-CM Codes	<p><b>H65.20</b> Chronic serous otitis media, unspecified ear</p> <p><b>H65.21</b> Chronic serous otitis media, right ear</p> <p><b>H65.22</b> Chronic serous otitis media, left ear</p> <p><b>H65.23</b> Chronic serous otitis media, bilateral</p> <p>* Use additional code for any associated perforated tympanic membrane (<b>H72-</b>)</p> <p>* Use additional code to identify:</p> <p>Exposure to environmental tobacco smoke (<b>Z77.22</b>)</p> <p>Exposure to tobacco smoke in the perinatal period (<b>P96.81</b>)</p> <p>History of tobacco use (<b>Z87.891</b>)</p> <p>Occupational exposure to environmental tobacco smoke (<b>Z57.31</b>)</p> <p>Tobacco dependence (<b>F17-</b>)</p> <p>Tobacco use (<b>Z72.0</b>)</p>
<b>470</b>	Deviated nasal septum <b>J34.2</b> Deviated nasal septum
<b>327.23</b>	Obstructive sleep apnea (adult)(pediatric)
ICD-10-CM Codes	<p><b>G47.33</b> Obstructive sleep apnea (adult)(pediatric)</p> <p>** Code also any associated underlying condition</p>

ICD-9-CM Code	ICD-10-CM Code(s)
<b>384.20</b>	Perforation of tympanic membrane, unspecified
ICD-10-CM Codes	<p><b>H72.90</b> Unspecified perforation of tympanic membrane, unspecified ear</p> <p>** Code first any associated otitis media (<b>H65-</b>, <b>H66.1-</b>, <b>H66.2-</b>, <b>H66.3-</b>, <b>H66.4-</b>, <b>H66.9-</b>, <b>H67-</b>)</p> <p><b>H72.91</b> Unspecified perforation of tympanic membrane, right ear</p> <p><b>H72.92</b> Unspecified perforation of tympanic membrane, left ear</p> <p><b>H72.93</b> Unspecified perforation of tympanic membrane, bilateral</p> <p>** There are more specific code choice selections available in ICD-10-CM. These include:</p> <p><b>H72.00</b> Central perforation of tympanic membrane, unspecified ear</p> <p><b>H72.01</b> Central perforation of tympanic membrane, right ear</p> <p><b>H72.02</b> Central perforation of tympanic membrane, left ear</p> <p><b>H72.03</b> Central perforation of tympanic membrane, bilateral</p> <p><b>H72.10</b> Attic perforation of tympanic membrane, unspecified ear</p> <p><b>H72.11</b> Attic perforation of tympanic membrane, right ear</p> <p><b>H72.12</b> Attic perforation of tympanic membrane, left ear</p> <p><b>H72.13</b> Attic perforation of tympanic membrane, bilateral</p> <p><b>H72.2x1</b> Other marginal perforations of tympanic membrane, right ear</p> <p><b>H72.2x2</b> Other marginal perforations of tympanic membrane, left ear</p> <p><b>H72.2x3</b> Other marginal perforations of tympanic membrane, bilateral</p> <p><b>H72.2x9</b> Other marginal perforations of tympanic membrane, unspecified ear</p> <p><b>H72.811</b> Multiple perforations of tympanic membrane, right ear</p> <p><b>H72.812</b> Multiple perforations of tympanic membrane, left ear</p> <p><b>H72.813</b> Multiple perforations of tympanic membrane, bilateral</p> <p><b>H72.819</b> Multiple perforations of tympanic membrane, unspecified ear</p> <p><b>H72.821</b> Total perforations of tympanic membrane, right ear</p> <p><b>H72.822</b> Total perforations of tympanic membrane, left ear</p> <p><b>H72.823</b> Total perforations of tympanic membrane, bilateral</p> <p><b>H72.829</b> Total perforations of tympanic membrane, unspecified ear</p>
<b>382.9</b>	Otitis media, unspecified
ICD-10-CM Codes	<p><b>H66.90</b> Otitis media, unspecified, unspecified ear</p> <p>* Use additional code for any associated perforated tympanic membrane (<b>H72-</b>)</p> <p>* Use additional code to identify:</p> <p>Exposure to environmental tobacco smoke (<b>Z77.22</b>)</p> <p>Exposure to tobacco smoke in the perinatal period (<b>P96.81</b>)</p> <p>History of tobacco use (<b>Z87.891</b>)</p> <p>Occupational exposure to environmental tobacco smoke (<b>Z57.31</b>)</p> <p>Tobacco dependence (<b>F17-</b>)</p> <p>Tobacco use (<b>Z72.0</b>)</p> <p>** There are more specific code choice selections available in ICD-10-CM. These include:</p> <p><b>H66.91</b> Otitis media, unspecified, right ear</p> <p><b>H66.92</b> Otitis media, unspecified, left ear</p> <p><b>H66.93</b> Otitis media, unspecified, bilateral</p> <p>** There are more specific codes available for otitis media. These include:</p> <p><b>H65.00</b> Acute serous otitis media, unspecified ear</p> <p><b>H65.01</b> Acute serous otitis media, right ear</p> <p><b>H65.02</b> Acute serous otitis media, left ear</p> <p><b>H65.03</b> Acute serous otitis media, bilateral</p> <p><b>H65.04</b> Acute serous otitis media, recurrent, right ear</p>



ICD-9-CM Code	ICD-10-CM Code(s)
<b>382.9</b>	Continued
ICD-10-CM Codes (cont.)	<p><b>H65.05</b> Acute serous otitis media, recurrent, left ear</p> <p><b>H65.06</b> Acute serous otitis media, recurrent, bilateral</p> <p><b>H65.07</b> Acute serous otitis media, recurrent, unspecified ear</p> <p><b>H65.111</b> Acute and subacute allergic otitis media (mucoïd) (sanguinous) (serous), right ear</p> <p><b>H65.112</b> Acute and subacute allergic otitis media (mucoïd) (sanguinous) (serous), left ear</p> <p><b>H65.113</b> Acute and subacute allergic otitis media (mucoïd) (sanguinous) (serous), bilateral</p> <p><b>H65.114</b> Acute and subacute allergic otitis media (mucoïd) (sanguinous) (serous), recurrent, right ear</p> <p><b>H65.115</b> Acute and subacute allergic otitis media (mucoïd) (sanguinous) (serous), recurrent, left ear</p> <p><b>H65.116</b> Acute and subacute allergic otitis media (mucoïd) (sanguinous) (serous), recurrent, bilateral</p> <p><b>H65.117</b> Acute and subacute allergic otitis media (mucoïd) (sanguinous) (serous), recurrent, unspecified ear</p> <p><b>H65.119</b> Acute and subacute allergic otitis media (mucoïd) (sanguinous) (serous), unspecified ear</p> <p><b>H65.191</b> Other acute nonsuppurative otitis media, right ear</p> <p><b>H65.192</b> Other acute nonsuppurative otitis media, left ear</p> <p><b>H65.193</b> Other acute nonsuppurative otitis media, bilateral</p> <p><b>H65.194</b> Other acute nonsuppurative otitis media, recurrent, right ear</p> <p><b>H65.195</b> Other acute nonsuppurative otitis media, recurrent, left ear</p> <p><b>H65.196</b> Other acute nonsuppurative otitis media, recurrent, bilateral</p> <p><b>H65.197</b> Other acute nonsuppurative otitis media, recurrent, unspecified ear</p> <p><b>H65.199</b> Other acute nonsuppurative otitis media, unspecified ear</p> <p><b>H65.20</b> Chronic serous otitis media, unspecified ear</p> <p><b>H65.21</b> Chronic serous otitis media, right ear</p> <p><b>H65.22</b> Chronic serous otitis media, left ear</p> <p><b>H65.23</b> Chronic serous otitis media, bilateral</p> <p><b>H65.30</b> Chronic mucoïd otitis media, unspecified ear</p> <p><b>H65.31</b> Chronic mucoïd otitis media, right ear</p> <p><b>H65.32</b> Chronic mucoïd otitis media, left ear</p> <p><b>H65.33</b> Chronic mucoïd otitis media, bilateral</p> <p><b>H65.411</b> Chronic allergic otitis media, right ear</p> <p><b>H65.412</b> Chronic allergic otitis media, left ear</p> <p><b>H65.413</b> Chronic allergic otitis media, bilateral</p> <p><b>H65.419</b> Chronic allergic otitis media, unspecified ear</p> <p><b>H65.491</b> Other chronic nonsuppurative otitis media, right ear</p> <p><b>H65.492</b> Other chronic nonsuppurative otitis media, left ear</p> <p><b>H65.493</b> Other chronic nonsuppurative otitis media, bilateral</p> <p><b>H65.499</b> Other chronic nonsuppurative otitis media, unspecified ear</p> <p><b>H65.90</b> Unspecified nonsuppurative otitis media, unspecified ear</p> <p><b>H65.91</b> Unspecified nonsuppurative otitis media, right ear</p> <p><b>H65.92</b> Unspecified nonsuppurative otitis media, left ear</p> <p><b>H65.93</b> Unspecified nonsuppurative otitis media, bilateral</p> <p><b>H66.001</b> Acute suppurative otitis media without spontaneous rupture of ear drum, right ear</p> <p><b>H66.002</b> Acute suppurative otitis media without spontaneous rupture of ear drum, left ear</p> <p><b>H66.003</b> Acute suppurative otitis media without spontaneous rupture of ear drum, bilateral</p> <p><b>H66.004</b> Acute suppurative otitis media without spontaneous rupture of ear drum, recurrent, right ear</p>

ICD-9-CM Code	ICD-10-CM Code(s)
ICD-10-CM Codes (cont.)	<p><b>H66.005</b> Acute suppurative otitis media without spontaneous rupture of ear drum, recurrent, left ear</p> <p><b>H66.006</b> Acute suppurative otitis media without spontaneous rupture of ear drum, recurrent, bilateral</p> <p><b>H66.007</b> Acute suppurative otitis media without spontaneous rupture of ear drum, recurrent, unspecified ear</p> <p><b>H66.009</b> Acute suppurative otitis media without spontaneous rupture of ear drum, unspecified ear</p> <p><b>H66.011</b> Acute suppurative otitis media with spontaneous rupture of ear drum, right ear</p> <p><b>H66.012</b> Acute suppurative otitis media with spontaneous rupture of ear drum, left ear</p> <p><b>H66.013</b> Acute suppurative otitis media with spontaneous rupture of ear drum, bilateral</p> <p><b>H66.014</b> Acute suppurative otitis media with spontaneous rupture of ear drum, recurrent, right ear</p> <p><b>H66.015</b> Acute suppurative otitis media with spontaneous rupture of ear drum, recurrent, left ear</p> <p><b>H66.016</b> Acute suppurative otitis media with spontaneous rupture of ear drum, recurrent, bilateral</p> <p><b>H66.017</b> Acute suppurative otitis media with spontaneous rupture of ear drum, recurrent, unspecified ear</p> <p><b>H66.019</b> Acute suppurative otitis media with spontaneous rupture of ear drum, unspecified ear</p> <p><b>H66.10</b> Chronic tubotympanic suppurative otitis media, unspecified</p> <p><b>H66.11</b> Chronic tubotympanic suppurative otitis media, right ear</p> <p><b>H66.12</b> Chronic tubotympanic suppurative otitis media, left ear</p> <p><b>H66.13</b> Chronic tubotympanic suppurative otitis media, bilateral</p> <p><b>H66.20</b> Chronic atticofacial suppurative otitis media, unspecified ear</p> <p><b>H66.21</b> Chronic atticofacial suppurative otitis media, right ear</p> <p><b>H66.22</b> Chronic atticofacial suppurative otitis media, left ear</p> <p><b>H66.23</b> Chronic atticofacial suppurative otitis media, bilateral</p> <p><b>H66.3X1</b> Other chronic suppurative otitis media, right ear</p> <p><b>H66.3X2</b> Other chronic suppurative otitis media, left ear</p> <p><b>H66.3X3</b> Other chronic suppurative otitis media, bilateral</p> <p><b>H66.3X9</b> Other chronic suppurative otitis media, unspecified ear</p> <p><b>H66.40</b> Suppurative otitis media, unspecified, unspecified ear</p> <p><b>H66.41</b> Suppurative otitis media, unspecified, right ear</p> <p><b>H66.42</b> Suppurative otitis media, unspecified, left ear</p> <p><b>H66.43</b> Suppurative otitis media, unspecified, bilateral</p> <p><b>H66.91</b> Unspecified otitis media, right ear</p> <p><b>H66.92</b> Unspecified otitis media, left ear</p> <p><b>H66.93</b> Unspecified otitis media, bilateral</p>
<b>474.01</b>	Chronic adenoiditis <b>J35.02</b> Chronic adenoiditis
<b>381.20</b>	Chronic mucoïd otitis media, simple or unspecified
ICD-10-CM Codes	<p><b>H65.30</b> Chronic mucoïd otitis media, unspecified ear</p> <p><b>H65.31</b> Chronic mucoïd otitis media, right ear</p> <p><b>H65.32</b> Chronic mucoïd otitis media, left ear</p> <p><b>H65.33</b> Chronic mucoïd otitis media, bilateral</p>



ICD-9-CM Code	ICD-10-CM Code(s)	
<b>473.9</b>	Unspecified sinusitis (chronic)	
ICD-10-CM Codes	<b>J32.9</b> Chronic sinusitis, unspecified * Use additional code to identify: Exposure to environmental tobacco smoke ( <b>Z77.22</b> ) Exposure to tobacco smoke in the perinatal period ( <b>P96.81</b> ) History of tobacco use ( <b>Z87.891</b> ) Occupational exposure to environmental tobacco smoke ( <b>Z57.31</b> ) Tobacco dependence ( <b>F17-</b> ) Tobacco use ( <b>Z72.0</b> ) ** There are more specific code choice selections available in ICD-10-CM. These include: <b>J32.0</b> Chronic maxillary sinusitis <b>J32.1</b> Chronic frontal sinusitis <b>J32.2</b> Chronic ethmoidal sinusitis <b>J32.3</b> Chronic sphenoidal sinusitis <b>J32.4</b> Chronic pansinusitis <b>J32.8</b> Other chronic sinusitis	
<b>784.7</b>	Epistaxis	<b>R04.0</b> Epistaxis
<b>475</b>	Peritonsillar abscess	<b>J36</b> Peritonsillar abscess
<b>473.8</b>	Chronic sinusitis, NEC	
ICD-10-CM Codes	<b>J32.8</b> Other chronic sinusitis * Use additional code to identify: Exposure to environmental tobacco smoke ( <b>Z77.22</b> ) Exposure to tobacco smoke in the perinatal period ( <b>P96.81</b> ) History of tobacco use ( <b>Z87.891</b> ) Occupational exposure to environmental tobacco smoke ( <b>Z57.31</b> ) Tobacco dependence ( <b>F17-</b> ) Tobacco use ( <b>Z72.0</b> ) ** There are more specific code choice selections available in ICD-10-CM. These include: <b>J32.0</b> Chronic maxillary sinusitis <b>J32.1</b> Chronic frontal sinusitis <b>J32.2</b> Chronic ethmoidal sinusitis <b>J32.3</b> Chronic sphenoidal sinusitis <b>J32.4</b> Chronic pansinusitis <b>J32.9</b> Chronic sinusitis, unspecified	
<b>210.2</b>	Benign neoplasm of major salivary glands	
ICD-10-CM Codes	<b>D11.0</b> Benign neoplasm of parotid gland <b>D11.9</b> Benign neoplasm of major salivary gland, unspecified <b>D11.7</b> Benign neoplasm of other major salivary glands	
<b>237.4</b>	Neoplasm of uncertain behavior of other and unspecified endocrine glands	
ICD-10-CM Codes	<b>D44.0</b> Neoplasm of uncertain behavior of thyroid gland <b>D44.2</b> Neoplasm of uncertain behavior of parathyroid gland <b>D44.9</b> Neoplasm of uncertain behavior of unspecified endocrine gland	
<b>239.1</b>	Neoplasm of respiratory system, unspecified nature	<b>D49.1</b> Neoplasm of unspecified behavior of respiratory system
<b>478.0</b>	Hypertrophy of nasal turbinates	<b>J34.3</b> Hypertrophy of nasal turbinates

ICD-9-CM Code	ICD-10-CM Code(s)	
<b>173.30</b>	Unspecified malignant neoplasm of skin of other and unspecified parts of face	
ICD-10-CM Codes	<b>C44.30</b> Malignant neoplasm of skin of unspecified part of face <b>C44.31</b> Malignant neoplasm of skin of nose <b>C44.39</b> Malignant neoplasm of skin of other parts of face <b>C44.300</b> Unspecified malignant neoplasm of skin of unspecified part of face <b>C44.301</b> Unspecified malignant neoplasm of skin of nose <b>C44.309</b> Unspecified malignant neoplasm of skin of other parts of face <b>C44.310</b> Basal cell carcinoma of skin of unspecified part of face <b>C44.311</b> Basal cell carcinoma of skin of nose <b>C44.319</b> Basal cell carcinoma of skin of other parts of face <b>C44.320</b> Squamous cell carcinoma of skin of unspecified parts of face <b>C44.321</b> Squamous cell carcinoma of skin of nose <b>C44.329</b> Squamous cell carcinoma of skin of other parts of face <b>C44.390</b> Other specified malignant neoplasm of skin of unspecified parts of face <b>C44.391</b> Other specified malignant neoplasm of skin of nose <b>C44.399</b> Other specified malignant neoplasm of skin of other parts of face	
<b>161.9</b>	Malignant neoplasm of larynx, unspecified	
ICD-10-CM Codes	<b>C32.9</b> Malignant neoplasm of larynx, unspecified * Use additional code to identify: Alcohol abuse and dependence ( <b>F10-</b> ) Exposure to environmental tobacco smoke ( <b>Z77.22</b> ) Exposure to tobacco smoke in the perinatal period ( <b>P96.81</b> ) History of tobacco use ( <b>Z87.891</b> ) Occupational exposure to environmental tobacco smoke ( <b>Z57.31</b> ) Tobacco dependence ( <b>F17-</b> ) Tobacco use ( <b>Z72.0</b> ) ** There are more specific code choice selections available in ICD-10-CM. These include: <b>C32.0</b> Malignant neoplasm of glottis <b>C32.1</b> Malignant neoplasm of supraglottis <b>C32.2</b> Malignant neoplasm of subglottis <b>C32.3</b> Malignant neoplasm of laryngeal cartilage <b>C32.8</b> Malignant neoplasm of overlapping sites of larynx	
<b>474.12</b>	Hypertrophy of adenoids	<b>J35.2</b> Hypertrophy of adenoids
<b>239.0</b>	Neoplasm of digestive system, unspecified nature	<b>D49.0</b> Neoplasm of unspecified behavior of digestive system
<b>235.1</b>	Neoplasm of lip, oral cavity, and pharynx, uncertain behavior	
ICD-10-CM Codes	<b>D37.01</b> Neoplasm of uncertain behavior of lip <b>D37.02</b> Neoplasm of uncertain behavior of tongue <b>D37.04</b> Neoplasm of uncertain behavior of minor salivary glands <b>D37.05</b> Neoplasm of uncertain behavior of pharynx <b>D37.09</b> Neoplasm of uncertain behavior of other specified sites of the oral cavity	
<b>784.2</b>	Swelling, mass, or lump in head and neck	
ICD-10-CM Codes	<b>R22.0</b> Localized swelling, mass and lump, head <b>R22.1</b> Localized swelling, mass and lump, neck	



ICD-9-CM Code	ICD-10-CM Code(s)
478.5	Other diseases of vocal cords <b>J38.3</b> Other diseases of vocal cords
706.2	Sebaceous cyst <b>L72.1</b> Trichodermal cyst
241.0	Nontoxic uninodular goiter <b>L72.3</b> Sebaceous cyst
173.20	Malignant neoplasm of skin of ear and external auditory canal
ICD-10-CM Codes	<b>C44.201</b> Unspecified malignant neoplasm of skin of unspecified ear and external auricular canal <b>C44.202</b> Unspecified malignant neoplasm of skin of right ear and external auricular canal <b>C44.209</b> Unspecified malignant neoplasm of skin of left ear and external auricular canal <b>C44.211</b> Basal cell carcinoma of skin of unspecified ear and external auricular canal <b>C44.212</b> Basal cell carcinoma of skin of right ear and external auricular canal <b>C44.219</b> Basal cell carcinoma of skin of left ear and external auricular canal <b>C44.221</b> Squamous cell carcinoma of skin of unspecified ear and external auricular canal <b>C44.222</b> Squamous cell carcinoma of skin of right ear and external auricular canal <b>C44.229</b> Squamous cell carcinoma of skin of left ear and external auricular canal <b>C44.291</b> Other specified malignant neoplasm of unspecified ear and external auricular canal <b>C44.292</b> Other specified malignant neoplasm of skin of right ear and external auricular canal <b>C44.299</b> Other specified malignant neoplasm of skin of left ear and external auricular canal
474.11	Hypertrophy of tonsils
ICD-10-CM Codes	<b>J35.1</b> Hypertrophy of tonsils * Use additional code to identify: Exposure to environmental tobacco smoke ( <b>Z77.22</b> ) Exposure to tobacco smoke in the perinatal period ( <b>P96.81</b> ) History of tobacco use ( <b>Z87.891</b> ) Occupational exposure to environmental tobacco smoke ( <b>Z57.31</b> ) Tobacco dependence ( <b>F17-</b> ) Tobacco use ( <b>Z72.0</b> )
802.0	Fracture of nasal bones, closed
ICD-10-CM Codes	<b>S02.2xxA</b> Fracture of nasal bones, initial encounter for closed fracture <b>S02.2xxD</b> Fracture of nasal bones, subsequent encounter for fracture with routine healing <b>S02.2xxG</b> Fracture of nasal bones, subsequent encounter for fracture with delayed healing <b>S02.2xxK</b> Fracture of nasal bones, subsequent encounter for fracture with nonunion <b>S02.2xxS</b> Fracture of nasal bones, sequela
527.2	Sialoadenitis
ICD-10-CM Codes	<b>K11.20</b> Sialoadenitis, unspecified <b>K11.21</b> Acute sialoadenitis <b>K11.22</b> Acute recurrent sialoadenitis <b>K11.23</b> Chronic sialoadenitis

ICD-9-CM Code	ICD-10-CM Code(s)
385.30	Cholesteatoma, unspecified
ICD-10-CM Codes	<b>H71.90</b> Unspecified cholesteatoma, unspecified ear <b>H71.91</b> Unspecified cholesteatoma, right ear <b>H71.92</b> Unspecified cholesteatoma, left ear <b>H71.93</b> Unspecified cholesteatoma, bilateral ** There are more specific code choice selections available in ICD-10-CM. These include: <b>H71.00</b> Cholesteatoma of attic, unspecified ear <b>H71.01</b> Cholesteatoma of attic, right ear <b>H71.02</b> Cholesteatoma of attic, left ear <b>H71.03</b> Cholesteatoma of attic, bilateral <b>H71.10</b> Cholesteatoma of tympanum, unspecified ear <b>H71.11</b> Cholesteatoma of tympanum, right ear <b>H71.12</b> Cholesteatoma of tympanum, left ear <b>H71.13</b> Cholesteatoma of tympanum, bilateral <b>H71.20</b> Cholesteatoma of mastoid, unspecified ear <b>H71.21</b> Cholesteatoma of mastoid, right ear <b>H71.22</b> Cholesteatoma of mastoid, left ear <b>H71.23</b> Cholesteatoma of mastoid, bilateral <b>H60.40</b> Cholesteatoma of external ear, unspecified ear <b>H60.41</b> Cholesteatoma of right external ear <b>H60.42</b> Cholesteatoma of left external ear <b>H60.43</b> Cholesteatoma of external ear, bilateral
471.8	Other polyp of sinus <b>J33.8</b> Other polyp of sinus
473.2	Chronic ethmoidal sinusitis
ICD-10-CM Codes	<b>J32.2</b> Chronic ethmoidal sinusitis * Use additional code to identify: Exposure to environmental tobacco smoke ( <b>Z77.22</b> ) Exposure to tobacco smoke in the perinatal period ( <b>P96.81</b> ) History of tobacco use ( <b>Z87.891</b> ) Occupational exposure to environmental tobacco smoke ( <b>Z57.31</b> ) Tobacco dependence ( <b>F17-</b> ) Tobacco use ( <b>Z72.0</b> )
530.6	Diverticulum of esophagus, acquired
ICD-10-CM Codes	<b>K22.5</b> Diverticulum of esophagus, acquired * Use additional code to identify: Exposure to environmental tobacco smoke ( <b>Z77.22</b> ) Exposure to tobacco smoke in the perinatal period ( <b>P96.81</b> ) History of tobacco use ( <b>Z87.891</b> ) Occupational exposure to environmental tobacco smoke ( <b>Z57.31</b> ) Tobacco dependence ( <b>F17-</b> ) Tobacco use ( <b>Z72.0</b> )
380.21	Cholesteatoma of external ear
ICD-10-CM Codes	<b>H60.40</b> Cholesteatoma of external ear, unspecified ear <b>H60.41</b> Cholesteatoma of right external ear <b>H60.42</b> Cholesteatoma of left external ear <b>H60.43</b> Cholesteatoma of external ear, bilateral
385.83	Retained foreign body of middle ear
ICD-10-CM Codes	<b>H74.8x1</b> Other specified disorders of right middle ear and mastoid <b>H74.8x2</b> Other specified disorders of left middle ear and mastoid <b>H74.8x3</b> Other specified disorders of middle ear and mastoid, bilateral <b>H74.8x9</b> Other specified disorders of middle ear and mastoid, unspecified ear



ICD-9-CM Code	ICD-10-CM Code(s)	
<b>173.40</b>	Malignant neoplasm of scalp and skin of neck	
ICD-10-CM Codes	<b>C44.4</b> Malignant neoplasm of skin of scalp and neck <b>C44.40</b> Unspecified malignant neoplasm of skin of scalp and neck <b>C44.41</b> Basal cell carcinoma of skin of scalp and neck <b>C44.42</b> Squamous cell carcinoma of skin of scalp and neck <b>C44.49</b> Other specified malignant neoplasm of skin of scalp and neck	
<b>239.7</b>	Neoplasm of endocrine glands and other parts of nervous system, unspecified nature	<b>D49.7</b> Neoplasm of unspecified behavior of endocrine glands and other parts of nervous system
<b>473.0</b>	Chronic maxillary sinusitis	<b>J32.0</b> Chronic maxillary sinusitis
<b>931</b>	Foreign body in ear	
ICD-10-CM Codes	<b>T16.1xxA</b> Foreign body in right ear, initial encounter <b>T16.1xxD</b> Foreign body in right ear, subsequent encounter <b>T16.1xxS</b> Foreign body in right ear, sequela <b>T16.2xxA</b> Foreign body in left ear, initial encounter <b>T16.2xxD</b> Foreign body in left ear, subsequent encounter <b>T16.2xxS</b> Foreign body in left ear, sequela <b>T16.9xxA</b> Foreign body in ear, unspecified ear, initial encounter <b>T16.9xxD</b> Foreign body in ear, unspecified ear, subsequent encounter <b>T16.9xxS</b> Foreign body in ear, unspecified ear, sequela	
<b>471.0</b>	Polyp of nasal cavity	
ICD-10-CM Codes	<b>J33.0</b> Polyp of nasal cavity * Use additional code to identify: Exposure to environmental tobacco smoke ( <b>Z77.22</b> ) Exposure to tobacco smoke in the perinatal period ( <b>P96.81</b> ) History of tobacco use ( <b>Z87.891</b> ) Occupational exposure to environmental tobacco smoke ( <b>Z57.31</b> ) Tobacco dependence ( <b>F17-</b> ) Tobacco use ( <b>Z72.0</b> )	
<b>478.4</b>	Polyp of vocal cord or larynx	
ICD-10-CM Codes	<b>J38.1</b> Polyp of vocal cord and larynx * Use additional code to identify: Exposure to environmental tobacco smoke ( <b>Z77.22</b> ) Exposure to tobacco smoke in the perinatal period ( <b>P96.81</b> ) History of tobacco use ( <b>Z87.891</b> ) Occupational exposure to environmental tobacco smoke ( <b>Z57.31</b> ) Tobacco dependence ( <b>F17-</b> ) Tobacco use ( <b>Z72.0</b> )	
<b>478.26</b>	Cyst of pharynx or nasopharynx	
ICD-10-CM Codes	<b>J39.0</b> Retropharyngeal and parapharyngeal abscess <b>J39.1</b> Other abscess of pharynx <b>J39.2</b> Other diseases of pharynx	

ICD-9-CM Code	ICD-10-CM Code(s)	
<b>193</b>	Malignant neoplasm of thyroid gland	
ICD-10-CM Codes	<b>C73</b> Malignant neoplasm of thyroid gland **Use additional code to identify any functional activity	
<b>195.0</b>	Malignant neoplasm of head, face, and neck	<b>C76.0</b> Malignant neoplasm of head, face, and neck
<b>141.9</b>	Malignant neoplasm of tongue, unspecified	
ICD-10-CM Codes	<b>C02.9</b> Malignant neoplasm of tongue, unspecified ** There are more specific code choice selections available in ICD-10-CM. These include: <b>C02.0</b> Malignant neoplasm of dorsal surface of tongue <b>C02.1</b> Malignant neoplasm of border of tongue <b>C02.2</b> Malignant neoplasm of ventral surface of tongue <b>C02.3</b> Malignant neoplasm of anterior two-thirds of tongue, part unspecified <b>C02.4</b> Malignant neoplasm of lingual tonsil <b>C02.8</b> Malignant neoplasm of overlapping sites of tongue	
<b>212.1</b>	Benign neoplasm of larynx	<b>D14.1</b> Benign neoplasm of larynx
<b>235.0</b>	Neoplasm of uncertain behavior of the major salivary glands	
ICD-10-CM Codes	<b>D37.030</b> Neoplasm of uncertain behavior of the parotid salivary glands <b>D37.031</b> Neoplasm of uncertain behavior of the sublingual salivary glands <b>D37.032</b> Neoplasm of uncertain behavior of the submandibular salivary glands <b>D37.039</b> Neoplasm of uncertain behavior of the major salivary glands, unspecified	
<b>787.20</b>	Dysphagia, unspecified	
ICD-10-CM Codes	<b>R13.10</b> Dysphagia, unspecified ** There are more specific code choice selections available in ICD-10-CM. These include: <b>R13.11</b> Dysphagia, oral phase <b>R13.12</b> Dysphagia, oropharyngeal phase <b>R13.13</b> Dysphagia, pharyngeal phase <b>R13.14</b> Dysphagia, pharyngoesophageal phase <b>R13.19</b> Other dysphagia	

*Fast Forward*

*to ICD-10*



ICD-10